SEMINARIO DE ANÁLISIS Y APLICACIONES

Lunes, 4 de mayo de 2020

12:00 h., ONLINE - URL: https://conecta.csic.es/b/jos-ajp-qp4

Glenier Bello

UAM-ICMAT

Models of linear operators satisfying operator inequalities

Resumen:

A general spectral theory of a linear operator does not exist, it only exists for particular subclasses of operators. One of the most celebrated theories of this type is the Nagy-Foiaş spectral theory of Hilbert space contractions, obtained in the 1960ies. It is based on the construction of a functional model, which heavily relies on certain chapters of Complex Analysis, such as the theory of Hardy spaces.

Being a contraction on a Hilbert space is characterized by a very simple operator inequality. In a landmark work of 1982, Agler showed how to pass from some other operator inequalities to a functional model of an operator by applying more general reproducing kernel Hilbert spaces instead of Hardy spaces. This work motivated an extensive research, which has become a rapidly growing branch of Operator Theory.

I will explain how to construct an explicit functional model for an operator satisfying a rather general operator inequality, and discuss the uniqueness of this model. Some spectral consequences in the spirit of the Nagy-Foiaş theory will be derived. I will also explain a new connection with the ergodic theory of linear operators.

Presentación previa a la defensa de la tesis doctoral.

ICMAT CSIC-UAM-UC3M-UCM Departamento de Matemáticas. U.A.M.



Universidad Autónoma de Madrid

