

Recovery of the Dirichlet-to-Neumann map from scattering data in the plane

By

KARI ASTALA*, DANIEL FARACO** and KEITH M. ROGERS***

Abstract

The differences between plane waves with fixed frequency and their analogues that have been distorted by a potential V yield the *scattering amplitude*. It is a classical problem to recover V from this information. It is well-known that the scattering amplitude uniquely determines the *Dirichlet-to-Neumann (DN) map* (from which the potential can be recovered) and there are a number of different approaches to proving this. Here we provide explicit formulae, closely following the work of Nachman and Stefanov, which recover the DN map from the scattering amplitude in the plane.

§ 1. Introduction

Let $\Omega \subset \mathbb{R}^2$ be a bounded domain which contains the support of a real potential V . We suppose throughout that $V \in L^p(\Omega)$, with $p > 2$, and that $k^2 > 0$ is not a Dirichlet eigenvalue for the Hamiltonian $-\Delta + V$. Then the Dirichlet-to-Neumann (DN) map Λ_V can be formally defined by

$$\Lambda_V : u|_{\partial\Omega} \mapsto \nabla u \cdot n|_{\partial\Omega},$$

where u is the solution to the Schrödinger equation

$$(1.1) \quad (-\Delta + V)u = k^2 u$$

Received April 20, 201x. Revised September 11, 201x.

2000 Mathematics Subject Classification(s): 2000 Mathematics Subject Classification(s): Primary 35P25, 45Q05; Secondary 35J10

Supported by the ERC grant 307179 and the MINECO grants MTM2010-16518, MTM2011-28198 and SEV-2011-0087 (Spain)

*Department of mathematics and statistics, Po.Box 68, 00014, University of Helsinki, Finland.

e-mail: kari.astala@helsinki.fi

**Departamento de Matemáticas - Universidad Autónoma de Madrid and Instituto de Ciencias Matemáticas CSIC-UAM-UC3M-UCM, 28049 Madrid, Spain.

e-mail: daniel.faraco@uam.es

***Instituto de Ciencias Matemáticas CSIC-UAM-UC3M-UCM, 28049 Madrid, Spain.

e-mail: keith.rogers@icmat.es

with the given Dirichlet data. In recent works (see for example [14, 11, 4, 2]), methods for recovering potentials from their DN maps have been developed.

On the other hand, for each $\theta \in \mathbb{S}^1$, the outgoing scattering solutions of (1.1) solve the Lippmann–Schwinger equation;

$$u(x, \theta) = e^{ikx \cdot \theta} - \int_{\mathbb{R}^2} G_0(x, y) V(y) u(y, \theta) dy.$$

Here, G_0 is the outgoing Green’s function which satisfies

$$(1.2) \quad (-\Delta - k^2)G_0(x, y) = \delta(x - y).$$

It can be calculated explicitly and is nothing more than a constant multiple of the zeroth order Hankel function of the first kind.

The scattering solutions satisfy the ‘asymptotics’ $u(x, \theta) = e^{ikx \cdot \theta}$ if $V = 0$. That is they measure how much the plane wave has been distorted by the potential. Indeed, later we will derive the asymptotics

$$G_0(x, y) = e^{-ik \frac{x}{|x|} \cdot y} \frac{e^{i\frac{\pi}{4}} e^{ik|x|}}{\sqrt{8\pi k|x|}} + o\left(\frac{1}{\sqrt{|x|}}\right),$$

so that, by plugging into the Lippmann–Schwinger equation, we obtain

$$u(x, \theta) = e^{ik\theta \cdot x} - A_V\left(\frac{x}{|x|}, \theta\right) \frac{e^{i\frac{\pi}{4}} e^{ik|x|}}{\sqrt{8\pi k|x|}} + o\left(\frac{1}{\sqrt{|x|}}\right),$$

Here, the scattering amplitude $A_V : \mathbb{S}^1 \times \mathbb{S}^1 \rightarrow \mathbb{C}$ satisfies

$$(1.3) \quad A_V(\vartheta, \theta) = \int_{\mathbb{R}^2} e^{-ik\vartheta \cdot y} V(y) u(y, \theta) dy,$$

and a classical problem is to recover the potential from this information alone.

It is well-known that the scattering amplitude uniquely determines the DN map (and *vice versa*) and so solutions to the first question, regarding the DN map, also provide solutions to the scattering question. There are a number of different approaches to showing this equivalence (see for example [3, 9, 16, 14, 15]). Here we provided explicit formulae, initially following an argument due to Nachman [10, Section 3] and then adapting three-dimensional arguments, due to Stefanov [13], to the two-dimensional problem.

§ 2. The formulae

It is well-known that, under the hypotheses of the introduction, for each $f \in H^{1/2}(\partial\Omega)$, there is a unique weak solution to the Dirichlet problem

$$(2.1) \quad \begin{cases} \Delta u = (V - k^2)u \\ u|_{\partial\Omega} = f, \end{cases}$$

that satisfies

$$(2.2) \quad \|u\|_{H^1(\Omega)} \leq C\|f\|_{H^{1/2}(\partial\Omega)}$$

(see for example [6] - in two dimensions $L^{n/2}(\mathbb{R}^n)$ can be replaced by $L^p(\mathbb{R}^2)$ with $p > 1$). Here $H^{1/2}(\partial\Omega) := H^1(\Omega)/H_0^1(\Omega)$, where $H_0^1(\Omega)$ denotes the closure of $C_0^\infty(\Omega)$ in $H^1(\Omega)$. The DN map Λ_V is then defined by

$$\langle \Lambda_V[f], \psi \rangle = \int_{\partial\Omega} \Lambda_V[f] \psi = \int_{\Omega} V u \Psi + \nabla u \cdot \nabla \Psi$$

for any $\Psi \in H^1(\Omega)$ with $\psi = \Psi + H_0^1(\Omega)$. When the solution and boundary are sufficiently regular, this definition coincides with that of the introduction by Green's formula. To see that Λ_V maps from $H^{1/2}(\partial\Omega)$ to $H^{-1/2}(\partial\Omega)$, the dual of $H^{1/2}(\partial\Omega)$, we note that by Hölder's inequality and the Hardy–Littlewood–Sobolev inequality,

$$\begin{aligned} \left| \langle \Lambda_V[f], \psi \rangle \right| &\leq \|u\|_{H^1(\Omega)} \|\Psi\|_{H^1(\Omega)} + \|V\|_p \|u\|_{L^q(\Omega)} \|\Psi\|_{L^q(\Omega)} \\ &\leq (1 + C\|V\|_p) \|u\|_{H^1(\Omega)} \|\Psi\|_{H^1(\Omega)} \end{aligned}$$

whenever $\Psi \in H^1(\Omega)$. Here $\frac{1}{p} + \frac{2}{q} = 1$ with $p > 1$. By (2.2), we obtain

$$\left| \langle \Lambda_V[f], \psi \rangle \right| \leq C(1 + \|V\|_p) \|f\|_{H^{1/2}(\partial\Omega)} \|\psi\|_{H^{1/2}(\partial\Omega)}$$

and so the DN map is bounded.

Essential to our analysis will be the outgoing Green's function G_V that satisfies

$$(2.3) \quad (-\Delta + V - k^2)G_V(x, y) = \delta(x - y)$$

and the corresponding near-field operator S_V defined via the single layer potential

$$S_V[f](x) = \int_{\partial\Omega} G_V(x, y) f(y) dy.$$

This is a bounded and invertible mapping from $H^{-1/2}(\partial\Omega)$ to $H^{1/2}(\partial\Omega)$ (the two-dimensional proof can be found in [7, Proposition A.1]). Then Nachman's formula [9],

$$\Lambda_V = \Lambda_0 + S_V^{-1} - S_0^{-1},$$

allows us to recover the DN map as soon as we recover the single layer potential S_V from the scattering amplitude A_V at energy k^2 . **change** For this, we also suppose that zero is not a Dirichlet eigenvalue of $-\Delta + V$ and that the boundary is sufficiently regular to apply Green's theorem.

When Ω is a disc, Nachman recovered S_V via formulae given by expansions in spherical harmonics as below. Otherwise he used a density argument (we remark that

Sylvester [15] also invokes density in order to recover). Since it is occasionally convenient to work with different domains (see for example [2] where it is convenient to work on a square), at this point we follow instead an argument of Stefanov [13], obtaining an explicit formula for the Green's function G_V in terms of A_V . Alternatively it seems likely that one could pass to the DN map on other domains from the the DN map on the disc via the argument in [11, Section 6] for the conductivity problem, however we prefer this more direct approach. We recover G_V outside of a disc which contains the potential, but which is properly contained in Ω , so that S_V can be obtained by integrating along the boundary $\partial\Omega$.

First we require the following well-known asymptotics.

Lemma 2.1. *Let $V \in L^p(\Omega)$ with $p > 2$. Then*

$$G_V(x, y) - G_0(x, y) = \frac{-i}{8\pi k} \frac{e^{ik|x|}}{|x|^{\frac{1}{2}}} \frac{e^{ik|y|}}{|y|^{\frac{1}{2}}} A_V\left(-\frac{x}{|x|}, \frac{y}{|y|}\right) + o\left(\frac{1}{|x|^{\frac{1}{2}}|y|^{\frac{1}{2}}}\right).$$

Proof. Using the asymptotics of the Hankel function for large r ;

$$H_0^{(1)}(r) = e^{-i\frac{\pi}{4}} \left(\frac{2}{\pi r}\right)^{\frac{1}{2}} e^{ir} + o\left(\frac{1}{r^{\frac{1}{2}}}\right)$$

(see for example [8, Section 5.16] or [5, pp. 66]) and the Taylor expansion at a fixed $y \in \mathbb{R}^2$,

$$|x - y| = |x| \left(1 - 2\frac{x}{|x|^2} \cdot y + \frac{|y|^2}{|x|^2}\right)^{1/2} = |x| - \frac{x}{|x|} \cdot y + O\left(\frac{1}{|x|}\right), \quad |x| \gg |y|,$$

one obtains the asymptotic formula

$$(2.4) \quad G_0(x, z) = \frac{i}{4} H_0^{(1)}(k|x - y|) = \frac{e^{i\frac{\pi}{4}}}{(8\pi)^{\frac{1}{2}}} \frac{e^{ik|x|}}{k^{\frac{1}{2}}|x|^{\frac{1}{2}}} e^{-ik\frac{x}{|x|} \cdot z} + o\left(\frac{1}{|x|^{\frac{1}{2}}}\right).$$

On the other hand, given that the outgoing solution to (2.3) is unique, one can verify that

$$(2.5) \quad G_V(x, y) = G_0(x, y) - \int_{\mathbb{R}^2} G_0(x, z) V(z) G_V(y, z) dz.$$

For this one must check that the outgoing Sommerfeld radiation condition is also uniformly satisfied for y in compact sets by the right-hand side. This follows from the fact that G_0 satisfies the condition (see for example [12, Proposition 2.1]) and that the resolvent

$$(-\Delta + V - k^2 - i0)^{-1} : V \mapsto \int_{\mathbb{R}^2} V(z) G_V(\cdot, z) dz$$

is bounded from $L^2((1 + |\cdot|^2)^\delta)$ to $L^2((1 + |\cdot|^2)^{-\delta})$ with $\delta > 1/2$ (see [1, Theorem 4.2]).

Similarly, the outgoing scattering solutions of (1.1), in the direction $-\frac{x}{|x|}$, solve

$$(2.6) \quad u(y, -\frac{x}{|x|}) = e^{-ik\frac{x}{|x|}\cdot y} - \int_{\mathbb{R}^2} e^{-ik\frac{x}{|x|}\cdot z} V(z) G_V(y, z) dy.$$

Plugging (2.4) into (2.5) and comparing with (2.6), we see that G_V satisfies

$$(2.7) \quad G_V(x, z) = \frac{e^{i\frac{\pi}{4}}}{(8\pi)^{\frac{1}{2}}} \frac{e^{ik|x|}}{k^{\frac{1}{2}}|x|^{\frac{1}{2}}} u\left(z, -\frac{x}{|x|}\right) + o\left(\frac{1}{|x|^{\frac{1}{2}}}\right).$$

As in (2.5), we also have that

$$(2.8) \quad G_V(x, y) - G_0(x, y) = - \int_{\mathbb{R}^2} G_0(y, z) V(z) G_V(x, z) dz.$$

Substituting (2.4) and (2.7) into this, we see that $G_V(x, y) - G_0(x, y)$ is equal to

$$\frac{-i}{8\pi k} \frac{e^{ik|x|}}{|x|^{\frac{1}{2}}} \frac{e^{ik|y|}}{|y|^{\frac{1}{2}}} \int_{\mathbb{R}^2} e^{-ik\frac{y}{|y|}\cdot z} V(z) u\left(z, -\frac{x}{|x|}\right) dz + o\left(\frac{1}{|x|^{\frac{1}{2}}|y|^{\frac{1}{2}}}\right),$$

so that, by using the formula (1.3), we obtain the result. \square

In the following theorem, $H_n^{(1)}$ denotes the Hankel function of the first kind and n th order (see for example [8] or [5]) and we write x in polar coordinates as $(|x|, \phi_x)$.

Theorem 2.2. *Let $V \in L^p(\Omega)$, with $p > 2$, be supported in the disc of radius ρ , centred at the origin, and consider its Fourier series*

$$A_V(\vartheta, \theta) = \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} a_{n,m} e^{in\phi_\vartheta} e^{im\phi_\theta}.$$

Then

$$G_V(x, y) - G_0(x, y) = \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} \frac{(-1)^n}{16} i^{n+m} a_{n,m} H_n^{(1)}(k|x|) H_m^{(1)}(k|y|) e^{in\phi_x} e^{im\phi_y},$$

where the series is uniformly and absolutely convergent for $|x| > |y| > R > \frac{3}{2}\rho > 0$.

Proof. We can expand $H_0^{(1)}(k|x - y|)$ as

$$H_0^{(1)}(k|x - y|) = H_0^{(1)}(k|x|) J_0(k|y|) + 2 \sum_{n \geq 1} H_n^{(1)}(k|x|) J_n(k|y|) \cos(\phi_x - \phi_y),$$

(see for example [5, Section 3.4] or [12, Theorem 3.4]). As $H_{-n}^{(1)} = (-1)^n H_n^{(1)}$ and $J_{-n} = (-1)^n J_n$, in order to separate variables it will be convenient to write this as

$$G_0(x, y) = \frac{i}{4} \sum_{n \in \mathbb{Z}} H_n^{(1)}(k|x|) J_n(k|y|) e^{in\phi_x} e^{-in\phi_y}.$$

Substituting (2.5) into (2.8) we obtain $G_V - G_0 = -I_1 + I_2$, where

$$I_1 = \int_{\mathbb{R}^2} G_0(x, z) V(z) G_0(z, y) dz$$

$$I_2 = \int_{\mathbb{R}^2} G_0(x, z_1) V(z_1) \int_{\mathbb{R}^2} G_V(z_1, z_2) V(z_2) G_0(y, z_2) dz_1 dz_2.$$

Now in both integrals we introduce the expansion of G_0 (recall that $G_0(z, y) = G_0(y, z)$), extracting the terms independent of z, z_1, z_2 . In this way we get

$$(2.9) \quad I_1 = \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} \alpha_{n,m} H_n^{(1)}(k|x|) H_m^{(1)}(k|y|) e^{in\phi_x} e^{im\phi_y},$$

$$(2.10) \quad I_2 = \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} \beta_{n,m} H_n^{(1)}(k|x|) H_m^{(1)}(k|y|) e^{in\phi_x} e^{im\phi_y},$$

where

$$\alpha_{n,m} = -\frac{1}{16} \int_{\mathbb{R}^2} V(z) J_n(k|z|) J_m(k|z|) e^{-i(n+m)\phi_z} dz,$$

$$\beta_{n,m} = -\frac{1}{16} \int_{\mathbb{R}^4} J_n(k|z_1|) V(z_1) G_V(z_1, z_2) V(z_2) J_m(k|z_2|) e^{-in\phi_{z_1}} e^{-im\phi_{z_2}} dz_1 dz_2.$$

It remains to show that the sums (2.9) and (2.10) converge uniformly and absolutely for $|x| > |y| > R > \frac{3}{2}\rho$. Once we know that this is the case, we can take limits and use the asymptotics of the Hankel functions for large r ;

$$H_n^{(1)}(r) = e^{-i(n\frac{\pi}{2} + \frac{\pi}{4})} \left(\frac{2}{\pi r}\right)^{\frac{1}{2}} e^{ir} + o\left(\frac{1}{r^{\frac{1}{2}}}\right)$$

(see for example [8, Section 5.16] or [5, pp. 66]), and then Lemma 2.1 tells us that

$$(-i)^{n+m+1} \frac{2}{\pi k} (\beta_{n,m} - \alpha_{n,m}) = -i \frac{(-1)^n}{8\pi k} a_{n,m}.$$

To see that the sums converge note that, by Hölder's inequality, we have

$$|\alpha_{n,m}| \leq C_\rho \|V\|_{L^p} \|J_n(k|\cdot|)\|_{L^\infty(B_\rho)} \|J_m(k|\cdot|)\|_{L^\infty(B_\rho)},$$

$$|\beta_{n,m}| \leq \|G_V\|_{L^2(B_\rho \times B_\rho)} \|V\|_{L^p}^2 \|J_n(k|\cdot|)\|_{L^\infty(B_\rho)} \|J_m(k|\cdot|)\|_{L^\infty(B_\rho)}.$$

At this point we deviate from [13] as there seems to be less local knowledge regarding G_V in two dimensions. Instead we can rewrite (2.8) as

$$G_V(\cdot, y) = G_0(\cdot, y) - (-\Delta + V - k^2 - i0)^{-1} [VG_0(\cdot, y)],$$

and use that the resolvent is bounded from $L^2((1 + |\cdot|^2)^\delta)$ to $L^2((1 + |\cdot|^2)^{-\delta})$ with $\delta > 1/2$ (see [1, Theorem 4.2]). Thus, using that V is compactly supported, and taking

$\frac{1}{2} = \frac{1}{p} + \frac{1}{q}$ with sufficiently large q ,

$$\begin{aligned} \|G_V(\cdot, y)\|_{L^2(B_\rho)} &\leq \|G_0(\cdot, y)\|_{L^2(B_\rho)} + C_\rho \|VG_0(\cdot, y)\|_{L^2(B_\rho)} \\ &\leq \|G_0(\cdot, y)\|_{L^2(B_\rho)} + C_\rho \|V\|_p \|G_0(\cdot, y)\|_{L^q(B_\rho)}. \end{aligned}$$

Integrating again with respect to y , and recalling that the singularity of $H_0^{(1)}$ at the origin is logarithmic, we see that $\|G_V\|_{L^2(B_\rho \times B_\rho)} \leq C_\rho(1 + \|V\|_p)$. Then, using the Taylor series expansion for the Bessel function,

$$|J_n(r)| = \left| \sum_{j \geq 0} \frac{(-1)^j}{j!(|n|+j)!} \left(\frac{r}{2}\right)^{2j+|n|} \right| \leq \frac{1}{|n|!} \left(\frac{\rho}{2}\right)^{|n|}$$

with $0 \leq r \leq \rho$, we see that

$$\begin{aligned} |\alpha_{n,m}| &\leq C_\rho \|V\|_p \frac{1}{|n|!} \left(\frac{k\rho}{2}\right)^{|n|} \frac{1}{|m|!} \left(\frac{k\rho}{2}\right)^{|m|}, \\ |\beta_{n,m}| &\leq C_\rho(1 + \|V\|_p)^3 \frac{1}{|n|!} \left(\frac{k\rho}{2}\right)^{|n|} \frac{1}{|m|!} \left(\frac{k\rho}{2}\right)^{|m|}. \end{aligned}$$

Plugging these estimates, along with the forthcoming estimate (2.11) for the Hankel functions, into the sums (2.9) and (2.10), we see that they are bounded by constant multiples of

$$\sum_{n \geq 0} \sum_{m \geq 0} \left(\frac{3\rho}{2R}\right)^n \left(\frac{3\rho}{2R}\right)^m$$

provided that $|x| > |y| > R > 0$. This series is of course convergent when $R > \frac{3}{2}\rho$, and so we are done. \square

For a fixed order n , the Hankel function of the first kind is well-known to decay at infinity as we saw earlier, however we need a bound that is uniform in n . The decay is not uniform in n as the singularity at the origin widens as n grows, however the following estimate is sufficient for our purposes.

Lemma 2.3. *For all $n \in \mathbb{Z}$ and all $r \geq R > 0$,*

$$(2.11) \quad |H_n^{(1)}(r)| \leq C_R |n|! \left(\frac{3}{R}\right)^{|n|}.$$

Proof. The Hankel functions have only one singularity, at the origin, and they decay at infinity (see for example [8, Section 5] or [5, Section 3.4]), so by taking the constant C_R sufficiently large, it will suffice to prove the estimate for $|n| \geq 2(R+6)$. As $H_{-n}^{(1)} = (-1)^n H_n^{(1)}$, we need only consider positive n and we divide these into two cases.

First we consider the easier case $4n(n-1) \leq r^2$ and use the recurrence relation

$$(2.12) \quad H_{n-1}^{(1)}(r) + H_{n+1}^{(1)}(r) = \frac{2n}{r} H_n^{(1)}(r)$$

(see for example [8, Section 5.4]) to conclude that

$$\begin{aligned} |H_{n+1}^{(1)}(r)| &= \left| \frac{2n}{r} \left(\frac{2(n-1)}{r} H_{n-1}^{(1)}(r) - H_{n-2}^{(1)}(r) \right) - H_{n-1}^{(1)}(r) \right| \\ &\leq |H_{n-1}^{(1)}(r)| + |H_{n-2}^{(1)}(r)|. \end{aligned}$$

Iterating this step, we see that

$$|H_n^{(1)}(r)| \leq 2^n \left(|H_0^{(1)}(r)| + |H_1^{(1)}(r)| + |H_2^{(1)}(r)| \right), \quad 4n(n-1) \leq r^2.$$

Taking $C_R = \sup_{r \geq R} (|H_0^{(1)}(r)| + |H_1^{(1)}(r)| + |H_2^{(1)}(r)|)$, and recalling that we can suppose that $n! \geq (2R/3)^n$ (by Stirling's formula $n > 2R$ is a stronger assumption), we obtain

$$(2.13) \quad |H_n^{(1)}(r)| \leq C_R n! \left(\frac{3}{R} \right)^n, \quad 4n(n-1) \leq r^2.$$

Next we consider the harder case $4n(n-1) > r^2$ and again use the recurrence relation (2.12) to conclude that

$$\begin{aligned} \frac{2n}{r} |H_n^{(1)}(r)| + |H_{n-1}^{(1)}(r)| &= \frac{2n}{r} \left| \frac{2(n-1)}{r} H_{n-1}^{(1)}(r) - H_{n-2}^{(1)}(r) \right| + |H_{n-1}^{(1)}(r)| \\ &\leq \frac{3n}{r} \left(\frac{2(n-1)}{r} |H_{n-1}^{(1)}(r)| + |H_{n-2}^{(1)}(r)| \right). \end{aligned}$$

Letting n_0 denote the smallest integer for which $4n_0(n_0-1) > r^2$, we iterate this inequality;

$$(2.14) \quad |H_{n+1}^{(1)}(r)| \leq \frac{3n}{r} \dots \frac{3n_0}{r} \left(\frac{2(n_0-1)}{r} |H_{n_0-1}^{(1)}(r)| + |H_{n_0-2}^{(1)}(r)| \right).$$

Now as long as $n_0 \geq 12$ (if $n_0 < 12$ then we are already done), we have

$$\frac{r}{2n_0} < \frac{2(n_0-1)}{r} \leq \frac{r}{2(n_0-2)} \leq \frac{6}{5} \frac{r}{2n_0},$$

which implies that $1/3 < 2(n_0-1)/r < 3$. For convenience we divide the first term and multiply the second term by this factor, so that

$$|H_{n+1}^{(1)}(r)| \leq 3 \frac{3n}{r} \dots \frac{3n_0}{r} \left(|H_{n_0-1}^{(1)}(r)| + \frac{2(n_0-1)}{r} |H_{n_0-2}^{(1)}(r)| \right).$$

Now as

$$\frac{3n}{r} \dots \frac{3n_0}{r} \leq \frac{3n}{R} \dots \frac{3n_0}{R} = n! \left(\frac{3}{R} \right)^n \frac{1}{(n_0-1)!} \left(\frac{R}{3} \right)^{n_0-1}$$

this yields

$$|H_{n+1}^{(1)}(r)| \leq 3n! \left(\frac{3}{R}\right)^n \left(\frac{1}{(n_0-1)!} \left(\frac{R}{3}\right)^{n_0-1} |H_{n_0-1}^{(1)}(r)| + \frac{1}{(n_0-2)!} \left(\frac{R}{3}\right)^{n_0-2} |H_{n_0-2}^{(1)}(r)| \right).$$

As n_0 was chosen to be the smallest integer belonging to the second case, we can bound $|H_{n_0-1}^{(1)}(r)|$ and $|H_{n_0-2}^{(1)}(r)|$ using the estimate (2.13) from the first case. This yields the estimate for the second case, and so the proof is complete. \square

The authors thank Juan Antonio Barceló, Adrian Nachman, Alberto Ruiz and Plamen Stefanov for helpful (electronic) conversations.

References

- [1] S. Agmon, Spectral properties of Schrödinger operators and scattering theory, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **2** (1975), no. 2, 151–218.
- [2] K. Astala, D. Faraco and K.M. Rogers, Rough potential recovery in plane, submitted.
- [3] J.M. Berezanskiĭ, The uniqueness theorem in the inverse problem of spectral analysis for the Schrödinger equation, Trudy Moskov. Mat. Obšč. **7** (1958), 1–62.
- [4] A.L. Bukhgeim, Recovering a potential from Cauchy data in the two-dimensional case, J. Inverse Ill-Posed Probl. **16** (2008), no. 1, 19–33.
- [5] D. Colton and R. Kress, *Inverse acoustic and electromagnetic scattering theory*, Applied Mathematical Sciences, 93, Springer, Berlin, 1992.
- [6] D. Dos Santos Ferreira, C. E. Kenig and M. Salo, Determining an unbounded potential from Cauchy data in admissible geometries, Comm. Partial Differential Equations **38** (2013), no. 1, 50–68.
- [7] V. Isakov and A. I. Nachman, Global uniqueness for a two-dimensional semilinear elliptic inverse problem, Trans. Amer. Math. Soc. **347** (1995), no. 9, 3375–3390.
- [8] N.N. Lebedev, *Special functions and their applications*, Revised English edition. Prentice Hall, Englewood Cliffs, NJ, 1965.
- [9] A.I. Nachman, Reconstructions from boundary measurements. Ann. of Math. **128** (1988), no. 3, 531–576.
- [10] A.I. Nachman, Inverse scattering at fixed energy, in *Mathematical physics, X (Leipzig, 1991)*, 434–441, Springer, Berlin. Nachman96, SU93a
- [11] A.I. Nachman, Global uniqueness for a two-dimensional inverse boundary value problem, Ann. of Math. **142** (1995), 71–96.
- [12] A. Ruiz, Harmonic analysis and inverse problems, www.uam.es/gruposinv/inversos/publicaciones/inverseproblems.pdf.
- [13] P. Stefanov, Stability of the inverse problem in potential scattering at fixed energy, Ann. Inst. Fourier (Grenoble) **40** (1990), no. 4, 867–884 (1991).
- [14] Z.Q. Sun and G. Uhlmann, Inverse scattering for singular potentials in two dimensions, Trans. Amer. Math. Soc. **338** (1993), no. 1, 363–374.
- [15] J. Sylvester, The Cauchy data and the scattering amplitude, Comm. Partial Differential Equations **19** (1994), no. 9-10, 1735–1741.
- [16] G. Uhlmann, Inverse boundary value problems and applications, Astérisque No. 207 (1992), 6, 153–211.