

ARC AND CURVE GRAPHS FOR INFINITE-TYPE SURFACES

JAVIER ARAMAYONA, ARIADNA FOSSAS & HUGO PARLIER

ABSTRACT. We study arc graphs and curve graphs for surfaces of infinite topological type. First, we define an arc graph relative to a finite number of (isolated) punctures and prove that it is a connected, uniformly hyperbolic graph of infinite diameter; this extends a recent result of J. Bavard to a large class of punctured surfaces.

Second, we study the subgraph of the curve graph spanned by those elements which intersect a fixed separating curve on the surface. We show that this graph has infinite diameter and geometric rank 3, and thus is not hyperbolic.

1. INTRODUCTION

For surfaces of finite topological type, an important number of problems about mapping class groups and Teichmüller spaces may be understood in terms of the various complexes constructed from curves and/or arcs. Prominent examples of these are the *curve graph* and the *arc graph* (see Section 3 for definitions); an important feature of both is that they are hyperbolic, see [7] and [9] respectively.

On the other hand, these complexes have received limited attention in the case of surfaces of infinite topological type, mainly due to the fact that mimicking the definitions from the case of finite-type surfaces ends up producing a graph of finite diameter; compare with Section 3. However, J. Bavard [1] has recently proved that a natural subgraph of the arc graph is hyperbolic and has infinite diameter (with respect to its intrinsic metric), in the case when the surface is homeomorphic to \mathbb{S}^2 minus the union of the north pole and a Cantor set. The main objective of this paper is to extend Bavard's result to a large class of punctured surfaces of infinite topological type.

1.1. Arc graphs. Let Σ be a connected orientable surface of infinite topological type and with empty boundary. Let $\Pi \subset \Sigma$ be the set of punctures of Σ , which we will always assume to be non-empty. It will also be useful to regard the elements of Π as marked points on Σ , and we will feel free to switch between the two viewpoints in the sequel. Throughout, we will need to assume that Π *contains at least one point that is isolated in Π* , when Π is equipped with the subspace topology.

Given a set P of isolated punctures, we define $\mathcal{A}(\Sigma, P)$ to be the simplicial graph whose vertices correspond to isotopy classes, relative to endpoints, of arcs on Σ with both endpoints in P , and where two such arcs are adjacent in $\mathcal{A}(\Sigma, P)$ if they can be realized disjointly on Σ ; see Section 3 for an expanded

definition. The graph $\mathcal{A}(\Sigma, P)$ turns into a metric space by deeming each edge to have unit length. We will prove:

Theorem 1.1. *Let Σ be a connected orientable surface, of infinite topological type and with at least one puncture. For any finite set of isolated P of punctures, the graph $\mathcal{A}(\Sigma, P)$ is connected, has infinite diameter, and is 7-hyperbolic.*

Remark. As mentioned above, in the particular case when Σ is homeomorphic to \mathbb{S}^2 minus the union of the north pole and a Cantor set, Theorem 1.1 is due to J. Bavard [1].

Theorem 1.1 could be regarded as a natural extension to the case of infinite-type surfaces of a result of Hensel-Przytycki-Webb [6], stated as Theorem 3.1 below, which asserts that arc graphs of finite-type surfaces are 7-hyperbolic. In fact, our proof relies heavily on this result.

However, it should be pointed out that, In spite of the similarities between the cases of finite- and infinite-type surfaces, the analogy between the two situations has limitations: indeed, a recent result of Bavard-Genevois [3] asserts that, unlike for finite-type surfaces, the mapping class group of an infinite-type surface is not *acylindrically hyperbolic*; see [10] for a definition.

A remark on the geometry of the different arc graphs. Consider the “full” arc graph $\mathcal{A}(\Sigma)$, whose vertices are *arbitrary* arcs with endpoints in Π , and where adjacency corresponds to disjointness. One easily sees that, as long as Π is infinite, the diameter of $\mathcal{A}(\Sigma)$ is equal to 2. This fact serves as justification for having to consider arc graphs *relative* to a finite set P of *isolated* punctures (as we will note in Section 3, the fact that the elements of P be isolated is essential to obtaining a graph of infinite diameter).

This said, the geometry of the different arcs graphs depends heavily on the subset of arcs used to define the given graph. On the one hand, the fact that arcs have *both* endpoints in P versus having *at least one* endpoint in P is unimportant, as the two definitions produce graphs that are quasi-isometric; see Lemma 3.4 below. However, in sharp contrast J. Bavard [?] has proved that the subgraph of $\mathcal{A}(\Sigma, P)$ spanned by those arcs that have *exactly one* endpoint in P is not hyperbolic whenever S has genus ≥ 1 or $|P| \geq 2$.

1.2. Curve graphs. Next, we turn our attention to curve graphs for surfaces of infinite topological type. Recall that, given a connected orientable surface Σ , the curve graph $\mathcal{C}(\Sigma)$ is the simplicial graph whose vertices are isotopy classes of essential simple closed curves on Σ , and where two such curves are adjacent in $\mathcal{C}(\Sigma)$ if they have disjoint representatives; see Section 3 for an expanded definition.

As was the case for arc graphs, when Σ has infinite topological type the graph $\mathcal{C}(\Sigma)$ has diameter 2 and thus has limited geometric interest. One natural way of producing a curve graph of infinite diameter is to only consider curves on Σ that intersect a fixed “portion” of Σ . More concretely, fix a separating curve α on Σ and consider the full subgraph $\mathcal{C}(\Sigma, \alpha)$ of $\mathcal{C}(\Sigma)$ spanned by those curves that essentially intersect α . Our second result states that $\mathcal{C}(\Sigma, \alpha)$ has infinite diameter but is never hyperbolic:

Theorem 1.2. *Let Σ be a connected orientable surface, of complexity ≥ 1 if Σ has finite topological type. If $\alpha \subset \Sigma$ is a separating curve, the graph $\mathcal{C}(\Sigma, \alpha)$ has geometric rank 3.*

Recall that the *geometric rank* of a metric space X is the largest integer n for which there is a quasi-isometric embedding of \mathbb{R}^n into X ; see Section 3 for an expanded definition.

We close the introduction by remarking that examples of non-hyperbolic complexes of infinite diameter were previously constructed by Fossas-Parlier [4]. Among these complexes is a type of pants graph for infinite type surfaces but, unlike the graphs we consider here, the graphs depend on the geometry of a fixed hyperbolic surface. However, they are very far from being hyperbolic as they have infinite geometric rank.

The plan of the paper is as follows. In Section 2 we will remind a few notions in metric geometry that will be used. Section 3 contains all the necessary background on arc and curve graphs. Section 4 deals with the proof of Theorem 1.1. Finally, in Section 5 we will give a proof of Theorem 1.2.

Acknowledgements. The first named author is supported by a Ramón y Cajal grant RYC-2013-13008. The second author is supported by ERC grant agreement number 267635 - RIGIDITY. The third author was supported by Swiss National Science Foundation grants numbers PP00P2_128557 and PP00P2_153024. The authors acknowledge support from U.S. National Science Foundation grants DMS 1107452, 1107263, 1107367 RNMS: Geometric structures And Representation varieties (the GEAR Network). We would like to thank Piotr Przytycki and Richard Webb for helpful conversations, and in particular for Proposition 3.3. We would also like to thank Julie Bavard for conversations, and in particular Proposition 3.5 and the first remark of Section 4.

2. PRELIMINARIES ON METRIC SPACES

We briefly recall the notions of Gromov hyperbolicity, quasi-isometry, and geometric rank. A nice discussion on these topics may be found in [5].

2.1. Hyperbolicity. Let (X, d) be a geodesic metric space, and $\delta \geq 0$. A geodesic triangle $T \subset X$ has a δ -center if there exists $c \in X$ whose distance to each of the three sides of T is at most δ . We say that (X, d) is δ -hyperbolic if every geodesic triangle in X has a δ -center; we will simply say that X is hyperbolic if it is δ -hyperbolic for some δ .

2.2. Quasi-isometries. Let (X, d_X) and (Y, d_Y) be two metric spaces. We say that a map $f : X \rightarrow Y$ is a (λ, C) -quasi-isometric embedding if there exist $\lambda \geq 1$ and $C \geq 0$ such that

$$\frac{1}{\lambda} \cdot d_X(x, y) - C \leq d_Y(f(x), f(y)) \leq \lambda \cdot d_X(x, y) + C$$

for all $x, y \in X$. We will say that the map f above is a *quasi-isometry* if, in addition, it is *almost surjective*: there exists $\delta > 0$ such that every element of Y is at distance at most δ from an element of $f(X)$.

We state the following observation as a separate lemma, as it is the way in which we will normally check that a given map is a quasi-isometry:

Lemma 2.1. *Let (X, d_X) and (Y, d_Y) be two metric spaces, and $f : X \rightarrow Y$ (resp. $g : Y \rightarrow X$) an L -Lipschitz (resp. K -Lipschitz) map. Suppose that there exist $D_f, D_g \geq 0$ such that $d(x, g \circ f(x)) \leq D_f$ and $d(y, f \circ g(y)) \leq D_g$ for all $x \in X$ and $y \in Y$. Then f (resp. g) is a quasi-isometry, with quasi-isometry constants that depend only on L and D_f (resp. K and D_g).*

Another well-known fact that we will heavily use is that *hyperbolicity is invariant under quasi-isometries*: if X is δ -hyperbolic and $f : X \rightarrow Y$ is a (λ, C) -quasi-isometry, then Y is δ' hyperbolic, where δ' depends (in an explicit way) only on δ , λ and C .

2.3. Geometric rank. As mentioned in the introduction, the *geometric rank* of a metric space X is the largest $n \in \mathbb{N}$ for which there exists a quasi-isometric embedding $\mathbb{R}^n \rightarrow X$. We note that an unbounded hyperbolic space has geometric rank 1; a nice exercise proves that the geometric rank is invariant under quasi-isometries.

3. ARC AND CURVE GRAPHS

In this section we define arc graphs and curve graphs, and state some results that will be used in the sequel. Throughout, Σ will be a connected orientable surface, possibly of infinite topological type, and with empty boundary. Recall that, in the case when Σ has finite topological type, the *complexity* of Σ is defined to be the number $3g - 3 + p$, where g and p are, respectively, the genus and number of punctures of Σ .

3.1. Arc graphs. Assume that Σ has at least one puncture, and let $\Pi \subset \Sigma$ be the set of punctures of Σ ; as mentioned in the introduction, we will often regard the elements of Π as marked points on Σ . By an arc on Σ we mean a nontrivial isotopy class (rel endpoints) of arcs on Σ , with endpoints in Π , and whose interior is disjoint from Π .

The *arc graph* $\mathcal{A}(\Sigma)$ is the simplicial graph whose vertices correspond to arcs on Σ , and where two arcs are adjacent in $\mathcal{A}(\Sigma)$ if they have representatives with disjoint interiors. The arc graph becomes a geodesic metric space by declaring the length of each edge to be 1. Masur-Schleimer proved that $\mathcal{A}(\Sigma)$ is δ -hyperbolic [9]; improving on this result, Hensel-Przytycki-Webb [6] recently proved that arc graphs of finite-type surfaces are *uniformly* hyperbolic:

Theorem 3.1 (Hensel-Przytycki-Webb [6]). *Let Σ be a surface of finite topological type, with at least one puncture. The arc graph $\mathcal{A}(\Sigma)$ is 7-hyperbolic.*

On the other hand, as we mentioned in the introduction the arc graph has little interest from a geometric point of view as long as Σ has infinitely many punctures, due to the following immediate observation:

Fact 3.2. *Let Σ be a connected orientable surface with an infinite number of punctures. Then $\mathcal{A}(\Sigma)$ has diameter 2.*

In order to overcome this obstacle, we will consider arc graphs *relative* to a finite set of punctures on Σ . More concretely, let $P \subset \Pi$ be a finite set, and consider the subgraph $\mathcal{A}(\Sigma, P) \subset \mathcal{A}(\Sigma)$ spanned by arcs on Σ with both endpoints in P . A minor adaptation of the arguments in [6] yields that, for Σ of finite topological type, these relative arc graphs are uniformly hyperbolic also, which will constitute a central ingredient of the proof of Theorem 1.1.

Proposition 3.3 (Hensel-Przytycki-Webb [6]). *Let Σ be a surface of finite topological type, and P a non-empty set of punctures of Σ . Then $\mathcal{A}(\Sigma, P)$ is 7-hyperbolic.*

Thus we see that our Theorem 1.1 is a natural extension of the above result to the context of infinite-type surfaces, provided these satisfy certain topological conditions.

Before we continue, recall from the introduction that choosing arcs to have both endpoints in P as opposed to having *at least* one endpoint in P is not important from the point of view of their large-scale geometry. More concretely, let P be a finite set of punctures on Σ whose every element is *isolated* in Π , equipped with the subspace topology. Consider the subgraph $\mathcal{A}^*(\Sigma, P)$ of $\mathcal{A}(\Sigma)$ spanned by those arcs with at least one endpoint in P . We have:

Lemma 3.4. *Let Σ be a connected orientable surface, of complexity at least 2 if it has finite topological type. The graphs $\mathcal{A}(\Sigma, P)$ and $\mathcal{A}^*(\Sigma, P)$ are quasi-isometric, with quasi-isometry constants that do not depend on Σ or P . In particular, for any finite-type surface Σ , $\mathcal{A}^*(\Sigma, P)$ is δ -hyperbolic for a universal constant δ that does not depend on Σ or P .*

Proof. We suppose that Σ has infinite topological type, as the finite-type case is easier. First, the natural inclusion map

$$\iota : \mathcal{A}(\Sigma, P) \rightarrow \mathcal{A}^*(\Sigma, P)$$

is 1-Lipschitz. In the other direction, there is a map

$$\phi : \mathcal{A}^*(\Sigma, P) \rightarrow \mathcal{A}(\Sigma, P)$$

defined as follows: given an arc $a \in \mathcal{A}^*(\Sigma, P)$, if $a \in \mathcal{A}(\Sigma, P)$ we define $\phi(a) = a$. Otherwise, suppose a has endpoints $p \in P$ and $q \notin P$. By the classification of infinite-type surfaces, the set of punctures of Σ is a subset of a Cantor set (see Proposition 5 of [11]). In the light of this, there exists a regular neighbourhood of a whose boundary does not contain any punctures. We define $\phi(a)$ to be the boundary of any such regular neighborhood, homotoped so that it is based at p (see Figure 1); here we are making use of the fact that p is isolated. We note that, while there may be many choices for $\phi(a)$, any two choices of $\phi(a)$ are at distance at most 2 in $\mathcal{A}(\Sigma, P)$.

Using the same reasoning, the images under the map ϕ of two disjoint arcs in $\mathcal{A}^*(\Sigma, P)$ are at distance at most 2 in $\mathcal{A}(\Sigma, P)$; in other words, ϕ is 2-Lipschitz.

Finally, observe that $a = \phi \circ \iota(a)$ for all $a \in \mathcal{A}(\Sigma, P)$. Similarly,

$$d(b, \iota \circ \phi(b)) \leq 1$$

for all $b \in \mathcal{A}^*(\Sigma, P)$ The result now follows from Lemma 2.1. \square




FIGURE 1. The image of the arc a under the map ϕ

Remark. Note that, a fortiori, the second assertion of Lemma 3.4 also holds for infinite-type surfaces, in the light of Theorem 1.1.

In sharp contrast, the words “at least” in the definition of $\mathcal{A}^*(\Sigma, P)$ are crucial, in the light of the following result due to J. Bavard [2].

Proposition 3.5 ([2]). *Let Σ be a surface of infinite topological type, and P a finite set of isolated punctures. Consider the subgraph $\mathcal{A}_*(\Sigma, P)$ of $\mathcal{A}(\Sigma, P)$ spanned by those arcs with exactly one endpoint in P . If Σ has genus at least 1, or if $|P| \geq 2$, then $\mathcal{A}_*(\Sigma, P)$ is not hyperbolic.*

3.2. Curve graphs. We say that a simple closed curve on Σ is *essential* if it does not bound a disk with at most one puncture. By a *curve* on Σ we will mean the free isotopy class of an essential simple closed curve on Σ .

The *curve graph* $\mathcal{C}(\Sigma)$ is the simplicial graph whose vertices are curves on Σ , and where two curves are adjacent in $\mathcal{C}(\Sigma)$ if they can be realized disjointly on Σ . As before, the curve graph turns into a geodesic metric space by deeming each edge to have length 1. A celebrated theorem of Masur-Minsky asserts that $\mathcal{C}(\Sigma)$ is hyperbolic whenever Σ has finite type:

Theorem 3.6 (Masur-Minsky [7]). *Let Σ be a connected orientable surface of finite topological type. If $\mathcal{C}(\Sigma)$ is connected then it is hyperbolic.*

As was the case with the arc graph, in the case of surfaces of infinite topological type the curve graph is not that interesting from a geometric viewpoint:

Fact 3.7. *Suppose Σ is a connected orientable surface of infinite topological type. Then $\mathcal{C}(\Sigma)$ has diameter 2.*

As mentioned in the introduction, one way of producing a “curve graph” of infinite diameter is to only consider curves on Σ that intersect a fixed curve on the surface. More concretely, fix a separating curve $\alpha \subset \Sigma$ and define $\mathcal{C}(\Sigma, \alpha)$ to be the full subgraph of $\mathcal{C}(\Sigma)$ spanned by those curves that essentially intersect α . However, in Theorem 1.2 we will prove that, while $\mathcal{C}(\Sigma, \alpha)$ has infinite diameter, it is never hyperbolic.

4. PROOF OF THEOREM 1.1

Throughout this section Σ will be a connected orientable surface of infinite topological type, with empty boundary, and with at least one puncture. Let Π be the set of punctures of Σ , and regard its elements as marked points

on Σ . We recall that, by the classification of infinite-type surfaces [11], the set Π is a subset of a Cantor set; see Proposition 5 of [11]. We will assume that Π contains at least one element that is isolated in Π equipped with the subspace topology. Let $P \subset \Pi$ a nonempty finite set of isolated punctures; we are going to show that $\mathcal{A}(\Sigma, P)$ is 7-hyperbolic.

Remark. If P contains a point that is not isolated, then $\mathcal{A}(\Sigma, P)$ has finite diameter. Indeed, the fact that arcs are compact implies that, for any two arcs with an endpoint on the same $p \in P$ not isolated, one can find a third arc based at p at distance at most 1 from both.

As we mentioned in the previous section, the main ingredient will be the result of Hensel-Przytycki-Webb [6] stated as Proposition 3.3 above. In order to be able to make use of their result, we need to define a certain type of *subsurface projection* map for arc graphs.

Subsurface projections:

Let $Y \subset \Sigma$ be a subsurface of finite topological type and complexity at least 2, with $P \subset Y$. Let $\mathcal{A}(Y, P)$ be the full subgraph of $\mathcal{A}(\Sigma, P)$ spanned by those vertices of $\mathcal{A}(\Sigma, P)$ that are entirely contained in Y . Fix, once and for all, an orientation on every boundary component of Y . We construct a map π_Y from $\mathcal{A}(\Sigma, P)$ to the power set of $\mathcal{A}(Y, P)$

$$\pi_Y : \mathcal{A}(\Sigma, P) \longrightarrow \mathcal{P}(\mathcal{A}(Y, P))$$

as follows:

- If c is entirely contained in Y , then $\pi_Y(c) := \{c\}$.
- Otherwise, consider any subarc c' of $c \subset Y$ that has one endpoint on $p \in P$. The other endpoint of c' necessarily lies on a boundary curve $\gamma \subset \partial Y$ (if not, c' would be entirely contained in Y). We orient c' so that it starts at p ; recall that γ has also been given an orientation. We define $\pi_Y(c')$ as the boundary of a regular neighborhood of $c' \circ \gamma \circ c'^{-1}$, homotoped so that it is based at p . We set

$$\pi_Y(c) := \{\pi_Y(c') \mid c' \text{ is a subarc of } c \text{ with one endpoint in } P\}.$$

In particular observe that, for any $c \in \mathcal{A}(\Sigma, P)$, the projection $\pi_Y(c)$ has at most two elements. The following lemma is an immediate consequence of the definition and will be key in the sequel.

Lemma 4.1. *Let $Y \subset \Sigma$ be a subsurface of finite topological type, of complexity at least 2, and with $P \subset Y$. Consider the map*

$$\pi_Y : \mathcal{A}(\Sigma, P) \rightarrow \mathcal{P}(\mathcal{A}(Y, P))$$

just defined. Then:

- (i) *For any vertex $c \in \mathcal{A}(\Sigma, P)$, we have*

$$\text{diam}_{\mathcal{A}(Y, P)}(\pi_Y(c)) \leq 2$$

- (ii) *Let $a, b \in \mathcal{A}(\Sigma, P)$ be disjoint arcs. Then, for any $c_a \in \pi_Y(a)$ and any $c_b \in \pi_Y(b)$, we have*

$$d_{\mathcal{A}(Y, P)}(c_a, c_b) \leq 2$$

Proof. To prove (i), let $c \in \mathcal{A}(\Sigma, P)$ and consider $d_1 \neq d_2 \in \pi_Y(c)$, with d_i coming from a subarc c'_i of c_i , $i = 1, 2$. Observe that c'_1 and c'_2 have disjoint interiors and, since $d_1 \neq d_2$, each c'_i has one endpoint in P and the other one in a boundary component $\gamma_i \subset \partial Y$. If $\gamma_1 \neq \gamma_2$ then $d_{\mathcal{A}(Y, P)}(d_1, d_2) = 1$. Otherwise d_1 and d_2 intersect twice and, since Y has complexity at least 2, there exists an arc in Y that is disjoint from both d_1 and d_2 . In particular, $d_{\mathcal{A}(Y, P)}(d_1, d_2) = 2$, as desired.

Part (ii) follows from the proof of part (i), since a and b are disjoint. \square

Since $\mathcal{A}(Y, P) \subset \mathcal{A}(\Sigma, P)$, we immediately obtain the following corollary of Lemma 4.1:

Corollary 4.2. *Let $Y \subset \Sigma$ be a subsurface of finite topological type, of complexity at least 2 and with $P \subset Y$. The inclusion map*

$$\mathcal{A}(Y, P) \hookrightarrow \mathcal{A}(\Sigma, P)$$

is a quasi-isometric embedding. More precisely, given $a, b \in \mathcal{A}(Y, P)$

$$d_{\mathcal{A}(\Sigma, P)}(a, b) \leq d_{\mathcal{A}(Y, P)}(a, b) \leq 2 d_{\mathcal{A}(\Sigma, P)}(a, b).$$

Proof. The left inequality comes from the fact that $\mathcal{A}(Y, P) \subset \mathcal{A}(\Sigma, P)$. To see that the right inequality holds, let $a, b \in \mathcal{A}(Y, P)$ and consider a path $[a, b]$ from a to b in $\mathcal{A}(\Sigma, P)$ between them. The projected path $\pi_Y([a, b])$ is a path from $\pi_Y(a) = a$ to $\pi_Y(b) = b$ in $\mathcal{A}(Y, P)$ which, by Lemma 4.1, has length at most twice that of $[a, b]$. Hence the result follows. \square

We need one more result before giving a proof of Theorem 1.1:

Lemma 4.3. *Let $F \subset \mathcal{A}(\Sigma, P)$ be a finite set of arcs. Then there exists a connected finite-type subsurface $Y \subset \Sigma$, of complexity at least 2 and with $P \subset Y$, such that every element of F is entirely contained in Y .*

Proof. First, we may enlarge F into a bigger finite set F' with the following two properties:

- (i) Every pair of distinct elements of P is the set of endpoints of an arc in F' .
- (ii) For any two pairs (p_1, q_1) and (p_2, q_2) of distinct elements of P , there exist arcs $c_1, c_2 \in F'$ such that c_i has endpoints in (p_i, q_i) , and c_1 and c_2 intersect.

As Π is a subset of a Cantor set and F' is finite, there exists a regular neighborhood Y of the union of the elements of F' that is a finite-type surface. Moreover, up to a further (finite) enlargement of F , we may assume that Y has complexity at least 2. We have $P \subset Y$ by (i); moreover, Y is connected by (ii). Since $F \subset F'$, it follows that every element of F is entirely contained in Y . \square

We are finally ready to prove Theorem 1.1:

Proof of Theorem 1.1. We first prove that $\mathcal{A}(\Sigma, P)$ is connected. Let $a, b \in \mathcal{A}(\Sigma, P)$. Let $Y \subset \Sigma$ be the subsurface given by applying Lemma 4.3 to the set $F = \{a, b\}$, so that we may view a, b as vertices of $\mathcal{A}(Y, P)$. Since

$\mathcal{A}(Y, P)$ is connected, there exists a path from a to b in $\mathcal{A}(Y, P)$ which, since $\mathcal{A}(Y, P) \subset \mathcal{A}(\Sigma, P)$, gives the desired path between a and b in $\mathcal{A}(\Sigma, P)$.

To see that $\mathcal{A}(\Sigma, P)$ has infinite diameter, choose a finite-type surface $Y \subset \Sigma$ of complexity at least 2 and with $P \subset Y$. Since $\mathcal{A}(Y, P)$ has infinite diameter, and is quasi-isometrically embedded in $\mathcal{A}(\Sigma, P)$ by Proposition 4.2, it follows that the diameter of $\mathcal{A}(\Sigma, P)$ is infinite.

We finally prove that $\mathcal{A}(\Sigma, P)$ is δ -hyperbolic for a universal constant δ ; in fact, as mentioned in the introduction, in this particular case $\mathcal{A}(\Sigma, P)$ will be 7-hyperbolic. Let $T \subset \mathcal{A}(\Sigma, P)$ be a geodesic triangle, and let F be the finite subset of $\mathcal{A}(\Sigma, P)$ whose elements are the vertices of T . Let Y be the connected, finite-type subsurface of Σ yielded by applying Lemma 4.3 to the set F ; in particular, we may view every vertex of T as an element of $\mathcal{A}(Y, P)$. Since $\mathcal{A}(Y, P)$ is 7-hyperbolic, by Proposition 3.3, it has a 7-center c , which is an arc in $\mathcal{A}(Y, P) \subset \mathcal{A}(\Sigma, P)$. Now the distances in $\mathcal{A}(\Sigma, P)$ from c to the sides of T are also at most 7. Since T is arbitrary, it follows that $\mathcal{A}(\Sigma, P)$ is 7-hyperbolic, as claimed. \square

5. PROOF OF THEOREM 1.2

In this section we will give a proof of Theorem 1.2. The strategy will be to prove that the graph $\mathcal{C}(\Sigma, \alpha)$ is quasi-isometric to a product of (three) hyperbolic spaces, namely arc graphs of subsurfaces determined by α .

Let Σ be a connected orientable surface, of complexity ≥ 1 if it has finite topological type, and fix a separating curve $\alpha \subset \Sigma$. Let Y_α be a closed regular neighborhood of α and denote by Y_1, Y_2 the connected components $\Sigma \setminus Y_\alpha$, so that $\Sigma = Y_1 \cup Y_2 \cup Y_\alpha$. Observe that Y_i has one more puncture than Y ; denote this new puncture by p_i .

Let $\mathcal{A}(Y_i, p_i)$ be the simplicial graph whose vertices are those isotopy classes of arcs in Y_i with both endpoints in p_i , and edges correspond to pairs of such arcs with disjoint representatives. Similarly, consider the simplicial graph $\mathcal{A}(Y_\alpha)$ whose vertices are isotopy classes, relative to ∂Y_α , of arcs in Y_α which connect the two boundary components of Y_α , and where edges correspond to pairs of such arcs with disjoint interiors. We note that, with respect to its natural path metric, $\mathcal{A}(Y_\alpha)$ is quasi-isometric to \mathbb{Z} , see Section 2.4 of [8] for a proof. Theorem 1.2 will be a consequence of the following:

Proposition 5.1. *The graph $\mathcal{C}(\Sigma, \alpha)$ is quasi-isometric to the product*

$$\mathcal{A}(Y_1, p_1) \times \mathcal{A}(Y_2, p_2) \times \mathcal{A}(Y_\alpha).$$

The claim above implies Theorem 1.2, since $\mathcal{A}(Y_1, p_1) \times \mathcal{A}(Y_2, p_2) \times \mathcal{A}(Y_\alpha)$ has geometric rank 3, being a product of three hyperbolic spaces.

Proof. For $i = 1, 2$, we define a map

$$\pi_i : \mathcal{C}(\Sigma, \alpha) \rightarrow \mathcal{P}(\mathcal{A}(Y_i, p_i))$$

given by $\pi_i(\beta) = \beta \cap Y_i$. Similarly, we set

$$\pi_\alpha : \mathcal{C}(\Sigma, \alpha) \rightarrow \mathcal{P}(\mathcal{A}(Y_\alpha))$$

informally by $\pi_\alpha(\beta) = \beta \cap Y_\alpha$; see Masur-Minsky's *projection to annuli* for a careful description of this map [8]. In this way we have constructed a map

$$\pi = (\pi_1, \pi_2, \pi_\alpha) : \mathcal{C}(\Sigma, \alpha) \rightarrow \mathcal{A}(Y_1, p_1) \times \mathcal{A}(Y_2, p_2) \times \mathcal{A}(Y_\alpha).$$

(Roughly speaking, the map π can be seen as a natural way of decomposing every curve β essentially intersecting α by a certain power of the Dehn twist along α and its behaviour away from α .) Since disjoint curves on $\mathcal{C}(\Sigma, \alpha)$ project to disjoint arcs on Y_i and Y_α , respectively, we deduce that $\pi_i(\beta)$ and $\pi_\alpha(\beta)$ are sets of diameter 1 in $\mathcal{A}(Y_i, p_i)$ and $\mathcal{A}(Y_\alpha)$, respectively. For the same reason, using an argument analogous to that of Proposition 4.2, we deduce that the maps π_i and π_α are 1-Lipschitz, and therefore we have:

Fact. The map π is 3-Lipschitz: for every $\beta, \gamma \in \mathcal{C}(\Sigma, \alpha)$ we have

$$d(\pi(\beta), \pi(\gamma)) \leq 3 \cdot d(\beta, \gamma).$$




FIGURE 2. Gluing arcs into a simple closed curve.

We now prove that there exist $\lambda \geq 1$ and $C \geq 0$ such that

$$(1) \quad \frac{1}{\lambda} \cdot d(\beta, \gamma) - C \leq d(\pi(\beta), \pi(\gamma)),$$

for all $\beta, \gamma \in \mathcal{C}(\Sigma, \alpha)$. In order to do so, we construct a map

$$\psi : \mathcal{A}(Y_1, p_1) \times \mathcal{A}(Y_2, p_2) \times \mathcal{A}(Y_\alpha) \rightarrow \mathcal{C}(\Sigma, \alpha)$$

that will serve as a coarse inverse of the map π , and we will apply Lemma 2.1. In order to construct this map it will be convenient to have an alternate description of the graphs $\mathcal{A}(Y_i, p_i)$, which we now give.

Let \bar{Y}_i be the two connected components of $\Sigma \setminus \text{int}(Y_\alpha)$, noting that $\bar{Y}_i = \text{int}(Y_i)$ for $i = 1, 2$. Let α_i be the boundary curve of \bar{Y}_i that is isotopic to α in Σ . Consider the simplicial graph $\mathcal{A}(\bar{Y}_i, \alpha_i)$ whose vertices are isotopy

classes of arcs on \overline{Y}_i with both endpoints on α_i , where isotopies need not fix α_i pointwise, and where two such arcs are adjacent in $\mathcal{A}(\overline{Y}_i, \alpha_i)$ if they can be realized disjointly. Observe that the graphs $\mathcal{A}(\overline{Y}_i, \alpha_i)$ and $\mathcal{A}(Y_i, p_i)$ are naturally isomorphic.

Armed with this alternate description, we proceed to construct the desired map ψ . Let $(a_1, a_2, b) \in \mathcal{A}(Y_1, p_1) \times \mathcal{A}(Y_2, p_2) \times \mathcal{A}(Y_\alpha)$, where we view the arc a_i as an element of $\mathcal{A}(\overline{Y}_i, \alpha_i)$ instead. We choose an arc $b' \in \mathcal{A}(Y_\alpha)$ that is disjoint from (but possibly equal to) b , and we glue the arcs a_1, b, b' and a_2 into a simple closed curve as shown in Figure 2. In this way we obtain the desired curve $\psi(a_1, a_2, b) \in \mathcal{C}(\Sigma, \alpha)$. We stress that while such curve is not unique, any two of them are within a uniformly bounded distance since they intersect a uniformly bounded number of times. For a totally analogous reason we deduce that there exists $L \geq 1$ such that ψ is L -Lipschitz, that is:

$$d(\psi(a_1, a_2, b), \psi(a'_1, a'_2, b')) \leq L \cdot d((a_1, a_2, b), (a'_1, a'_2, b'))$$

for all triples (a_1, a_2, b) and (a'_1, a'_2, b') . Moreover, it is immediate from the constructions of the maps π and ψ that there exists a universal constant $R > 0$, not depending on Σ or α , such that

$$d(\pi \circ \psi(a_1, a_2, b), (a_1, a_2, b)) \leq R$$

for all triples (a_1, a_2, b) . Similarly, there exists a constant $R' > 0$, again not depending on Σ or α , such that

$$d(\psi \circ \pi(\beta), \beta) \leq R'$$

for all $\beta \in \mathcal{C}(\Sigma, \alpha)$. At this point, the result follows from Lemma 2.1. \square

REFERENCES

- [1] J. Bavard, *Hyperbolicité du graphe des rayons et quasi-morphismes sur un gros groupe modulaire*, to appear in Geometry and Topology.
- [2] J. Bavard, *personal communication*.
- [3] J. Bavard, A. Genevois, *in preparation*.
- [4] A. Fossas, H. Parlier, *Curve graphs on surfaces of infinite type*, to appear in Ann. Acad. Sci. Fenn. Math.
- [5] E. Ghys, P. de la Harpe (editors), *Sur les groupes hyperboliques d'après Mikhael Gromov*. Papers from the Swiss Seminar on Hyperbolic Groups held in Bern, 1988. Progress in Mathematics, 83. Birkhuser Boston, Inc., Boston, MA, 1990.
- [6] S. Hensel, P. Przytycki, R. Webb, *1-slim triangles and uniform hyperbolicity for arc graphs and curve graphs*. J. Eur. Math. Soc. (JEMS) 17 (2015)
- [7] Masur, Minsky, *Geometry of the complex of curves. I. Hyperbolicity*. Invent. Math. 138 (1999)
- [8] Masur, Minsky, *Geometry of the complex of curves. II. Hierarchical structure*. Geom. Funct. Anal. 10 (2000)
- [9] H. Masur, S. Schleimer, *The geometry of the disk complex*, Journal of the American Mathematical Society, 26 (2013).
- [10] D. Osin, *Acylically hyperbolic groups*, Trans. Amer. Math. Soc. (2015).
- [11] I. Richards, *On the classification of noncompact surfaces*, Trans. Amer. Math. Soc. 106 (1963).